


EBBING • GAMMON

General  
**Chemistry**  
ELEVENTH EDITION

# Chemical Reactions

## 4.5 Oxidation–Reduction Reactions



Iron nail and copper(II) sulfate solution, which has a blue color.



Fe reacts with  $\text{Cu}^{2+}(aq)$  to yield  $\text{Fe}^{2+}(aq)$  and  $\text{Cu}(s)$ . The molecular view above the pictures depicts the net ionic equation; water and the sulfate anion have been omitted.



The copper metal forms a coating on the nail.





The net ionic equation is:



## ➤ Oxidation Numbers

an **oxidation–reduction reaction** (or **redox reaction**)  
is a reaction in which electrons are transferred between species or  
in which atoms change oxidation number.

Formerly, the term *oxidation* meant “reaction with oxygen.”



# ➤ Oxidation-Number Rules:

Table 4.5 Rules for Assigning Oxidation Numbers


Copyright 2017 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. WCN 02-200-203

Rule	Applies to	Statement
1	Elements	The oxidation number of an atom in an element is zero.
2	Monatomic ions	The oxidation number of an atom in a monatomic ion equals the charge on the ion.
3	Oxygen	The oxidation number of oxygen is $-2$ in most of its compounds. (An exception is O in $\text{H}_2\text{O}_2$ and other peroxides, where the oxidation number is $-1$ .)
4	Hydrogen	The oxidation number of hydrogen is $+1$ in most of its compounds. (The oxidation number of hydrogen is $-1$ in binary compounds with a metal, such as $\text{CaH}_2$ .)
5	Halogens	The oxidation number of fluorine is $-1$ in all of its compounds. Each of the other halogens (Cl, Br, I) has an oxidation number of $-1$ in binary compounds, except when the other element is another halogen above it in the periodic table or the other element is oxygen.
6	Compounds and ions	The sum of the oxidation numbers of the atoms in a compound is zero. The sum of the oxidation numbers of the atoms in a polyatomic ion equals the charge on the ion.


Examples:



# ➤ Describing Oxidation–Reduction Reactions





We can write this reaction in terms of two half-reactions



## ➤ Some Common Oxidation–Reduction Reactions

1. Combination reaction
2. Decomposition reaction
3. Displacement reaction
4. Combustion reaction



1. **Combination Reactions** is a reaction in which two substances combine to form a third substance



✓ Not all combination reactions are oxidation- reduction reactions



**2. Decomposition Reactions** is a reaction in which a single compound reacts to give two or more substances





✓ Not all decomposition reactions are oxidation-reduction reactions




**3. Displacement reaction** (also called a **single-replacement reaction**) is a reaction in which an element reacts with a compound, displacing another element from it.

✓ involve an element and one of its compounds → must be oxidation–reduction reactions.






Net ionic rxn.



- ✓ metals listed at the top are the strongest reducing agents (they lose electrons easily)
- ✓ A free element reacts with the monatomic ion of another element if the free element is above the other element in the activity series
- ✓ The highlighted elements react slowly with liquid water, but readily with steam, to give H<sub>2</sub>





**Table 4.6** Activity Series of the Elements

React vigorously with acidic solutions and water to give H <sub>2</sub>	$\left\{ \begin{array}{l} \text{Li} \\ \text{K} \\ \text{Ba} \\ \text{Ca} \\ \text{Na} \end{array} \right.$
React with acids to give H <sub>2</sub>	$\left\{ \begin{array}{l} \text{Mg} \\ \text{Al} \\ \text{Zn} \\ \text{Cr} \\ \text{Fe} \\ \text{Cd} \\ \text{Co} \\ \text{Ni} \\ \text{Sn} \\ \text{Pb} \end{array} \right.$
Do not react with acids to give H <sub>2</sub> *	$\left\{ \begin{array}{l} \text{H}_2 \\ \text{Cu} \\ \text{Hg} \\ \text{Ag} \\ \text{Au} \end{array} \right.$







**4. Combustion reaction** is *a reaction in which a substance reacts with oxygen, usually with the rapid release of heat to produce a flame.*

✓ The products include one or more oxides. Oxygen changes oxidation number from 0 to -2, so combustions are oxidation-reduction reactions.








## 4.6 Balancing Simple Oxidation-Reduction Equations

(Q) Apply the half-reaction method to balance the following equation:

$$\text{Mg}(s) + \text{N}_2(g) \rightarrow \text{Mg}_3\text{N}_2(s)$$


**4.66** Balance the following oxidation–reduction reactions by the half-reaction method.



## 4.7 Molar Concentration

$$\text{Molarity (}M\text{)} = \frac{\text{moles of solute}}{\text{liters of solution}}$$

(Q) A sample of  $\text{NaNO}_3$  weighing 0.38 g is placed in a 50.0 mL volumetric flask. The flask is then filled with water to the mark on the neck. What is the molarity of the resulting solution?

$$\text{Molarity} = \frac{4.47 \times 10^{-3} \text{ mol NaNO}_3}{50.0 \times 10^{-3} \text{ L soln}} = \mathbf{0.089 M \text{ NaNO}_3}$$

## 4.8 Diluting Solutions

$$M_i \times V_i = M_f \times V_f$$

(Q) You are given a solution of 14.8  $M$   $\text{NH}_3$ . How many milliliters of this solution do you require to give 100.0 mL of 1.00  $M$   $\text{NH}_3$  ?

$$V_i = \frac{1.00 \cancel{M} \times 100.0 \text{ mL}}{14.8 \cancel{M}} = \mathbf{6.76 \text{ mL}}$$

✓ Number of moles does not change

(Q) What is the molar concentration of  $\text{Na}^+$  in a solution made by dissolving 1.59 g of  $\text{Na}_2\text{CO}_3$  (molar mass = 106g/mol) in 100 mL  $\text{H}_2\text{O}$ ?

## 4.9 Gravimetric Analysis

*is a type of quantitative analysis in which the amount of a species in a material is determined by converting the species to a product that can be isolated completely and weighed.*

(Q) A 1.000-L sample of polluted water was analyzed for lead(II) ion,  $\text{Pb}_{2+}$ , by adding an excess of sodium sulfate to it. The mass of lead(II) sulfate that precipitated was 229.8 mg. What is the mass of lead in a liter of the water? Give the answer as milligrams of lead per liter of solution.

✓ Solution: mass percentage of Pb in  $\text{PbSO}_4$

$$\% \text{ Pb} = \frac{207.2 \text{ g/mol}}{303.3 \text{ g/mol}} \times 100\% = 68.32\%$$

Amount Pb in sample = 229.8 mg  $\text{PbSO}_4$  X 0.6832 = 157.0 mg Pb

The water sample contains **157.0 mg Pb per liter.**

Exercise 4.14 You are given a sample of limestone, which is mostly  $\text{CaCO}_3$ , to determine the mass percentage of Ca in the rock. You dissolve the limestone in hydrochloric acid, which gives a solution of calcium chloride. Then you precipitate the calcium ion in solution by adding sodium oxalate,  $\text{Na}_2\text{C}_2\text{O}_4$ . The precipitate is calcium oxalate,  $\text{CaC}_2\text{O}_4$ . You find that a sample of limestone weighing 128.3 mg gives 140.2 mg of  $\text{CaC}_2\text{O}_4$ . What is the mass percentage of calcium in the limestone?


**4.85** Copper has compounds with copper(I) ion or copper(II) ion. A compound of copper and chlorine was treated with a solution of silver nitrate,  $\text{AgNO}_3$ , to convert the chloride ion in the compound to a precipitate of  $\text{AgCl}$ . A 59.40-mg sample of the copper compound gave 86.00 mg  $\text{AgCl}$ .

- a. Calculate the percentage of chlorine in the copper compound.
- b. Decide whether the formula of the compound is  $\text{CuCl}$  or  $\text{CuCl}_2$ .

## 4.10 Volumetric Analysis

### Example 4.13 Calculating the Volume of Reactant Solution Needed

(Q) Consider the following reaction:



Suppose a beaker contains 35.0 mL of 0.175 M  $\text{H}_2\text{SO}_4$ . How many milliliters of 0.250 M NaOH must be added to react completely with the sulfuric acid?

$$35.0 \times 10^{-3} \text{ L } \cancel{\text{H}_2\text{SO}_4 \text{ soln}} \times \frac{0.175 \text{ mol } \cancel{\text{H}_2\text{SO}_4}}{1 \text{ L } \cancel{\text{H}_2\text{SO}_4 \text{ soln}}} \times \frac{2 \text{ mol } \cancel{\text{NaOH}}}{1 \text{ mol } \cancel{\text{H}_2\text{SO}_4}} \times \frac{1 \text{ L NaOH soln}}{0.250 \text{ mol } \cancel{\text{NaOH}}} = 4.90 \times 10^{-2} \text{ L NaOH soln (or 49.0 mL NaOH soln)}$$

Exercise 4.15 consider the following reaction:




How many milliliters of 0.375 M NiSO<sub>4</sub> will react with 45.7 mL of 0.265 M Na<sub>3</sub>PO<sub>4</sub>?

## Example 4.14

### Calculating the Quantity of Substance in a Titrated Solution

(Q) A flask contains a solution with an unknown amount of HCl. This solution is titrated with 0.207 M NaOH. It takes 4.47 mL of the NaOH solution to complete the reaction. What is the mass of the HCl?



**Solution** The calculation is as follows:

$$4.47 \times 10^{-3} \text{ L NaOH soln} \times \frac{0.207 \text{ mol NaOH}}{1 \text{ L NaOH soln}} \times \frac{1 \text{ mol HCl}}{1 \text{ mol NaOH}} \times \frac{36.5 \text{ g HCl}}{1 \text{ mol HCl}} = 0.0338 \text{ g HCl}$$


**4.91** How many milliliters of 0.150 M  $\text{H}_2\text{SO}_4$  are required to react with 8.20 g of  $\text{NaHCO}_3$ , according to the following equation?



**4.111** A stock solution of potassium dichromate,  $K_2Cr_2O_7$ , is made by dissolving 84.5 g of the compound in 1.00 L of solution. How many milliliters of this solution are required to prepare 1.00 L of 0.15 M  $K_2Cr_2O_7$ ?

**4.113** A solution contains 6.0% (by mass) NaBr. The density of the solution is 1.046 g/cm<sup>3</sup>. What is the molarity of NaBr?

**4.132** Identify each of the following reactions as being a neutralization, precipitation, or reduction-oxidation reaction.



**4.135**(modified) A 25-mL sample of 0.50 *M* NaOH is combined with a 75-mL sample of 0.30 *M* NaOH. What is the molarity of the resulting NaOH solution?

**4.140** Potassium hydrogen phthalate (abbreviated as KHP) has the molecular formula  $\text{KHC}_8\text{H}_4\text{O}_4$  and a molar mass of 204.22 g/mol. KHP has one acidic hydrogen. A solid sample of KHP is dissolved in 50 mL of water and titrated to the equivalence point with 22.90 mL of a 0.5010 M NaOH solution. How many grams of KHP were used in the titration?

**4.74** What is the volume (in milliliters) of 0.100 M  $\text{H}_2\text{SO}_4$  containing 0.949 g  $\text{H}_2\text{SO}_4$ ?