



The background features abstract, overlapping green geometric shapes in various shades, including light lime green, medium green, and dark forest green. These shapes are primarily located on the left and right sides of the slide, framing the central white area.

The Cell


Hanan Jafar. BDS.MSc.PhD



Cell polarity



- ▶ Many cells show polarity, meaning different areas of the cell have different structures
- ▶ The most-studied polarity is in epithelial cells, they have
 - ▶ Apical domain
 - ▶ Basal (basolateral) domain



The Plasma Membrane

The background of the slide features abstract, overlapping geometric shapes in various shades of green, ranging from light lime to dark forest green. These shapes are primarily located on the right side and bottom of the frame, creating a modern, layered effect against the white background.

What is the plasma membrane?

- ▶ The **plasma membrane** (cell membrane or plasmalemma) that envelops every eukaryotic cell.
- ▶ It functions as a selective barrier regulating the passage of materials into and out of the cell and facilitating the transport of specific molecules.

IMPORTANT



- ▶ Membranes range from 7.5 to 10 nm in thickness and consequently are visible only in the electron microscope (can NOT be resolved by the light microscope). They appear as a trilaminar unit in TEM.
- ▶ The line between adjacent cells sometimes seen faintly with the light microscope consists of plasma membrane proteins plus extracellular material, which together can reach a dimension visible by light microscopy.

Components of plasma membrane


- ▶ Phospholipids
- ▶ Cholesterol
- ▶ Proteins
 - ▶ Integral: incorporated directly within the lipid bilayer
 - ▶ Peripheral: bound to one of the two membrane surfaces, particularly on the cytoplasmic side
- ▶ Oligosaccharide (carbohydrate) chains linked to many of the phospholipid (to form glycolipids) and protein (to form glycoproteins) molecules.

Structure of plasma membrane


- Model: Mosaic phospholipid bilayer




b



a





b



Functions of Plasma Membrane

- 1. Physical barrier:** Establishes a flexible boundary, protects cellular contents, and supports cell structure. Phospholipid bilayer separates substances inside and outside the cell
- 2. Selective permeability:** Regulates entry and exit of ions, nutrients, and waste molecules through the plasma membrane
- 3. Electrochemical gradients:** Establishes and maintains an electrical charge difference across the plasma membrane
- 4. Communication:** Contains receptors that recognize and respond to molecular signals



The Cytoplasm


The background of the slide is white with abstract, overlapping green geometric shapes on the right side. These shapes include triangles and polygons in various shades of green, from light lime to dark forest green, creating a modern, layered effect.

Overview

- ▶ Inside the cell membrane, the fluid cytoplasm (or **cytosol**) bathes metabolically active structures called organelles, which may be membranous (such as mitochondria) or nonmembranous protein complexes (such as ribosomes).
- ▶ Most organelles are positioned in the cytoplasm by movements along the polymers of the cytoskeleton, which also determines a cell's shape and motility.

Ribosomes

- ▶ **Ribosomes** are macromolecular machines, about 20×30 nm in size, which assemble polypeptides (proteins) from amino acids in a sequence specified by mRNA.
- ▶ They can be free in the cytosol or bound to the rough endoplasmic reticulum
- ▶ During protein synthesis many ribosomes typically bind the same strand of mRNA to form larger complexes called **polyribosomes**, or **polysomes**




Differences between free and bound ribosomes

- ▶ Free ribosomes synthesize cytosolic and cytoskeletal proteins and proteins for import into the nucleus, mitochondria, and peroxisomes.
- ▶ Bound ribosomes synthesize proteins that are to be incorporated into membranes, stored in lysosomes, or eventually secreted from the cell.
 - ▶ The proteins produced by these ribosomes are segregated during translation into the interior of the ER's membrane cisternae.

Endoplasmic Reticulum (ER)

- ▶ The **endoplasmic reticulum** is an anastomosing network of intercommunicating channels or **cisternae** formed by a continuous membrane network.
- ▶ Of two types:
 - ▶ Rough endoplasmic reticulum (rER)
 - ▶ Smooth endoplasmic reticulum (sER)



Functions of ER

- ▶ **Synthesis:** Provides a place for chemical reactions
 - ▶ sER is the site of lipid synthesis and carbohydrate metabolism
 - ▶ rER synthesizes proteins for secretion, incorporation into the plasma, membrane, and as enzymes within lysosomes
- ▶ **Transport:** Moves molecules through cisternal space from one part of the cell to another, sequestered away from the cytoplasm
- ▶ **Storage:** Stores newly synthesized molecules, sER stores Ca^{++}
- ▶ **Detoxification:** sER detoxifies both drugs and alcohol




Histological appearance



▶ rER:



- ▶ Light microscopy: Intense basophilia (blue to purple on H & E)
- ▶ Electron microscopy: Appears as interconnected flat cisternae and tubules associated with ribosomes

▶ sER:

- ▶ Light microscopy: can not be seen under LM
- ▶ Electron microscopy: Appears as interconnected tubules with various shapes and sizes and not stack of flattened cisternae not associated with ribosomes








Smooth ER


Rough ER

0.2 μm




Golgi Apparatus

- ▶ **Golgi apparatus**, or Golgi complex, completes posttranslational modifications of proteins produced in the rER and then packages and addresses these proteins to their proper destinations.
- ▶ Material moves from the rER cisternae to the Golgi apparatus in small, membrane-enclosed carriers called **transport vesicles**
- ▶ Has two sides (ends):
 - ▶ Receiving end (cis): receives transport vesicles
 - ▶ Shipping end (tran): ships secretory vesicles





Histological appearance



- ▶ Golgi cannot be seen in H & E staining.
- ▶ In highly active cells, with prominent golgi apparatus, it gives a negative image (as if an empty space)



Secretory Granules



- ▶ The granules are surrounded by membrane and contain a concentrated form of the secretory product
- ▶ Histologically:
 - ▶ Light microscopy: they cannot be resolved, but in cells active in protein synthesis, they give apical cytoplasm intense eosinophilic appearance (pink)
 - ▶ Electron microscopy: homogenous electron dense structures near the apex of the cell







Lysosomes

- ▶ **Lysosomes** are sites of intracellular digestion and turnover of cellular components.
- ▶ Under electron microscopy, we can distinguish between primary and secondary lysosomes:
 - Primary: Uniformly granular electron dense appearance
 - Secondary: Larger with heterogenous appearance (particulate content)
- ▶ Lysosomes are not well shown on H&E-stained cells but can be visualized by light microscopy after staining with toluidine blue.




- ▶ Synthesis of lysosomal enzymes occurs in the RER, with packaging in the Golgi apparatus. Endocytosis produces vesicles that fuse with **endosomes** before merging with **lysosomes**.
- ▶ Phagocytic vacuoles (or phagosomes) fuse with primary lysosomes to become **secondary lysosomes** (or heterolysosomes), in which ingested material is degraded.
- ▶ **Autophagosomes**, such as those depicted here with a mitochondrion in the process of digestion, are formed after nonfunctional or surplus organelles become enclosed with membrane and the resulting structure fuses with a lysosome.
- ▶ The products of lysosomal digestion are recycled to the cytoplasm, but indigestible molecules remain in a membrane-enclosed **residual body**, which may accumulate in long-lived cells as lipofuscin.
- ▶ In some cells, such as osteoclasts, the lysosomal enzymes are secreted into a restricted extracellular compartment.



Mitochondria

- ▶ **Mitochondria** are membrane-enclosed organelles with arrays of enzymes specialized for aerobic respiration and production of **adenosine triphosphate (ATP)**, which supplies energy for most cellular activities.
- ▶ The number of mitochondria is related to the cell's energy needs: cells with a high-energy metabolism (eg, cardiac muscle, cells of some kidney tubules) have abundant mitochondria, whereas cells with a low-energy metabolism have few mitochondria.


- ▶ Under the TEM each mitochondrion is seen to have two separated and very different membranes that together create two compartments: the innermost **matrix** and a narrow **intermembrane space**
- ▶ The **outer membrane** contains many transmembrane proteins called **porins** that form channels through which small molecules such as pyruvate and other metabolites pass from the cytoplasm to the intermembrane space.
- ▶ The **inner membrane** has many long folds called **cristae**, which project into the matrix and greatly increase the membrane's surface area





The cytoskeleton

- ▶ The cytoplasmic **cytoskeleton** is a complex array of:
 - (1) microtubules,
 - (2) microfilaments (also called actin filaments)
 - (3) intermediate filaments.
- ▶ These protein polymers determine the shapes of cells, play an important role in the movements of organelles and cytoplasmic vesicles, and also allow the movement of entire cells.



General Function of Cytoskeleton




- Structural:** Provides structural support to cell; stabilizes junctions between cells
- Movement:** Assists with cytosol streaming and cell motility; helps move organelles and materials throughout cell; helps move chromosomes during cell division

Microtubules

Microfilaments

Intermediate filaments


Polymer



Centrosome

- ▶ The **centrosome** is the microtubule-organizing center for the mitotic spindle and consists of paired centrioles.
- ▶ The TEM reveals that the two centrioles in a centrosome exist at right angles to each other.
- ▶ Each centriole consists of **nine microtubular triplets** peripherally arranged, and no microtubules in the center (arrangement: 9+0)

Centrosome





The Nucleus

The background features abstract, overlapping geometric shapes in various shades of green, ranging from light lime to dark forest green. These shapes are primarily located on the right side of the frame, creating a modern, layered effect. The rest of the background is plain white.

The Nucleus

- ▶ The nucleus contains the code for all of a cell's enzymes and other proteins. It also contains the molecular machinery to replicate the DNA and to synthesize and process all types of RNA.
- ▶ The nucleus usually appears as a large rounded or oval structure, often near the cell's center.
- ▶ It consists of a nuclear envelope containing chromatin, with one or more specialized regions of chromatin called **nucleoli**.




Functions of the Nucleus

1. **Cellular regulation:** Houses genetic material, which directs all cellular activities and regulates cellular structure
2. **Production:** Produces ribosomal subunits in nucleolus and exports them into cytoplasm for assembly into ribosomes


Chromatin

- ▶ The mass of DNA and its associated proteins
- ▶ Microscopically two categories of chromatin can be distinguished:
 - ▶ **Euchromatin** is visible as finely dispersed granular material in the electron microscope and as lightly stained basophilic areas in the light microscope.
 - associated with active cells
 - ▶ **Heterochromatin** appears as coarse, electron-dense material in the electron microscope and as intensely basophilic clumps in the light microscope.
 - associated with inactive cells




Nucleolus

- ▶ The **nucleolus** is a spherical, highly basophilic subdomain of nuclei in cells actively engaged in protein synthesis
- ▶ It is the location of ribosomal subunit assembly and transcription of ribosomal RNA (**rRNA**)
- ▶ The intense basophilia of nucleoli is due **not** to heterochromatin but to the presence of densely concentrated **rRNA**



The Nuclear Envelope

- ▶ The **nuclear envelope** is a double set of membranes with a narrow perinuclear space, which separates the cytoplasm from nucleoplasm
- ▶ The outer membrane binds ribosomes and is continuous with the RER.
- ▶ It is penetrated by **nuclear pore complexes**,
- ▶ It is supported internally by a meshwork, the nuclear lamina, composed of intermediate filament subunits called **lamins**.



Nuclear pore complexes

- ▶ Nuclear pore complexes (nuclear pores) contain more than 30 core proteins (nucleoporins), span both membranes of the nuclear envelope, and regulate the bidirectional transfer of macromolecular complexes between the nucleus and cytoplasm

