


MEASURING DISEASE OCCURRENCE

INCIDENCE AND PREVALENCE (MORBIDITY MEASURES)

Dr. Sireen Alkhalidi, BDS, MPH, DrPH
Department of Family and Community
Medicine
School of Medicine/ The University of
Jordan
Second Semester 2020/2021



How do we measure diseases?

Four *quantitative* descriptors to measure disease occurrence:

- ▶ Numbers
- ▶ Ratios
- ▶ Proportions
- ▶ Rates

Descriptors

Numbers: Use of actual number of events
e.g 100 cases of TB in community A

Ratios: Quantifies the magnitude of one occurrence
X, in relation to another event Y as X/Y
e.g Ratio of TB cases in community A to B is 1:10

Descriptors

Proportions: a ratio in which the numerator is included in the denominator

e.g proportion of TB cases in community A is 10%

Rates: a proportion with time element

It measure the occurrence of an event overtime

e.g US measles cases in 2000/US population in 2000

Measurement of Disease Occurrence

Morbidity measures


5

Morbidity rates are rates that are used to quantify the magnitude/frequency of diseases

Two common morbidity measures:

Incidence rates (Cumulative incidence, incidence density)

Prevalence (Period prevalence, point prevalence)




Incidence rate

- ✓ The proportion of a population that develops a disease overtime
- ✓ The risk/probability of an individual developing a disease overtime
- ✓ The rapidity with which new cases of a disease develop overtime
- ✓ The proportion of unaffected individuals who on average will contract the disease overtime

Cumulative incidence

7

Cumulative Incidence = $\frac{\text{Number of new cases of a disease during a specified period}}{\text{Population at risk in the same Period of time}}$



Practical challenges in measuring incidence rate

1. Identification of population at risk


Population at risk constitutes all those free of the disease and susceptible to it

2. Population is not static/it fluctuates/as a result of births, deaths and migration

3. People are at risk only until they get the disease and then no more at risk

Prevalence

It measures the proportion of a population with a disease during a specified period or at a point in time



prevalence

Measures the proportion of a population with a disease at a point in time

$$\text{prevalence} = \frac{\text{All persons with a disease}}{\text{Total population}}$$

It is not a rate, but a true proportion

Incidence Vs prevalence

11

Prevalence measures all of the current cases of the disease in the community.


- ✓ It depends on the duration of the disease process
- ✓ It depends on the incidence of the disease
- ❖ It can be used to determine the health care needs of a community.

$$P = I \times D$$

where **P** = Prevalence rate, **I** = Incidence rate, **D** = Duration of the disease.

- ✓ **Prevalence rate is equal to Incidence rate in case of diseases with short duration or highly fatal such as Rabies (after onset of clinical symptoms).**

Prevalence vs. Incidence



Relationship between prevalence & incidence rates

13

An increase in prevalence may not necessarily be due to an increase in incidence rate, it could be due to an increase in average duration of a disease due to decrease in death and/or recovery rates.

Prevalence = Incidence Rate x Average Duration

If: the frequency of disease is rare (i.e., <10% of the population has it).

- ✓ If the average duration of disease remains constant, then preventive measures that reduce the incidence of disease would be expected to result in a decreased prevalence.
- ✓ Similarly, if the incidence remained constant, then developing a cure would reduce the average duration of disease, and this would also reduce the prevalence of disease.
- ✓ **In the late 1990s anti-retroviral therapy was introduced and greatly improved the survival of people with HIV. However, they weren't cured of their disease, meaning that the average duration of disease increased. As a result, the prevalence of HIV increased during this period.**

Prevalence = Incidence Rate x Average Duration

- ✓ The relationship can be visualized by thinking of inflow and outflow from a reservoir. The fullness of the reservoir can be thought of as analogous to prevalence, and Raindrops might represent incidence, or the rate at which new cases of a disease are being added to the population, thus becoming prevalent cases.
- ✓ Water also flows out of the reservoir, analogous to removal of prevalent cases by virtue of either dying or being cured of the disease.
- ✓ Imagine that incidence (rainfall) and the rate of cure or death are initially equal; if so, the height of water in the reservoir will remain constant.

Calculation ...

A survey of respiratory disease was conducted and the results are presented in the table below.

Calculate the prevalence of chronic bronchitis in each age group and in the total group.

Prevalence of chronic bronchitis, by age, in a sample of 2383 employed men: , 1981.			
Age (years)	Number Surveyed	Frequency	Prevalence (%)
45-49	496	18	3.6
50-54	672	18	2.7
55-59	1215	18	1.5
Total	2383	54	2.3
$\chi^2 = 0.983, p = 0.612$			

$$\begin{aligned}\text{Prevalence} &= 54 / 2383 = 0.0226 \times 100\% = 2.3\% \\ &= 0.0226 \times 1000 = 22.6 \text{ cases/ } 1000 \text{ pop.}\end{aligned}$$

A study was conducted to examine the incidence of Carpal Tunnel Syndrome (CTS) among computer operators in a certain corporation. An initial survey was given to 12 administrative assistants. Two of the 12 administrative assistants had symptoms and 10 did not reveal signs or symptoms equivalent to CTS. The administrative assistants who did not reveal signs or symptoms equivalent to CTS were then recruited into a study and followed for 4 years. The findings are listed below

3 of the 10 administrative assistants developed CTS during the 4 year follow-up period

Subjects	Follow-up Time(yrs)	CTS
1	1	yes
1	2.5	yes
1	3	yes
2	2	fired
1	1	transferred
4	4	no

Calculate Cumulative Incidence (per 1,000).

Cumulative Incidence= $3 / 10 = 0.3 \times 100\% = 30\%$

$= 0.3 \times 1000 = 300$ cases per 1,000 population