

# Cornea and Sclera

---



Muawyah Al Bdour

The University of Jordan

2023/2024




# Objectives

---

- Know the basic anatomy, histology, physiology
- Highlight some common diseases
- Special focus on Keratoconus being relatively common in the region
- Brief notes on corneal graft

# Cornea and sclera: tough outer coat



0.5 mm thick  
11-12 mm in diameter

## 5 Layers:

**Epithelium:** Non-keratinized stratified squamous. **Regenerates** from stem cells found in the limbus


## Bowman membrane

**Stroma:** 90 % of corneal thickness; parallel collagen fibrils


**Descemet membrane:** tough

**Endothelium:** a monolayer of **non-regenerating** cells that actively pump water from the stroma to control hydration of the cornea. When damaged, by disease or surgery, cells spread

Loss of barrier and pumping functions lead to edema and corneal clouding



- 
- ❑ The cornea derives its nutrition almost entirely from aqueous humour which supplies  $O_2$  to the endothelium & post. stroma. While the ant. stroma & epithelium receive  $O_2$  from ambient air and tear film
  - ❑ Avascular
  - ❑ Rich in sensory nerve endings originating from the nasociliary branch of trigeminal nerve



---

The focus must be adjustable to allow clear vision for both distant and near objects ( with accommodation).

## **Function of the cornea:**

- Protection of internal ocular structures
- Refraction

Refractive components of the eye or the focusing power is :

- \*\* 2/3 cornea, fixed power
- \*\* 1/3 lens, power increases with accommodation

# *Cornea*

---

**Infective corneal lesions**

# Infective corneal lesions


## Herpes Simplex and Zoster Keratitis

---

Latency of the virus in the trigeminal ganglion

- Recurrent infection involve reactivation of the latent virus
- The risk of reactivation increases in debilitated patients and with stress

- **HSV type (1) & (2), HZV (Varicella Zoster)**
- Primary infection
- **Recurrent infection** results from reactivation
- Characterized by the appearance of dendritic ulcers on the cornea, which **usually** heal without scar





- 
- HSV type 1 is the most common cause
  - Usually acquired in early life due close contact and **what we see in clinic is reactivation**
  - Symptoms and signs:
    - Very painful and associated with photophobia, watery discharge and swelling of the eyelids
    - Pathognomonic appearance is *dendritic ulcer*

# HZV


- Dendritic ulcers
- **Skin lesions:** start as vesicle at ophthalmic division of Trig
- Pain and headache
- Also need medical & derma care
- Ocular problems are more likely if the nasociliary branch is involved **Hutchinson's sign**

↳ think of it as the branch that supplies the tip of the nose passes first to give the cornea / so if it's involved → then most probably the cornea is also involved.  
(eye)



- 
- If the stroma is involved ( disciform keratitis which is immunogenic reaction to herpes antigen) > corneal edema > permanent scarring > corneal graft may be required

- **Rx:**

- Topical antiviral for dendritic ulcer ( acyclovir )
- ~~Topical steroids must be avoided in patients with dendritic ulcer~~ since they may cause more extensive ulceration 
- Oral antiviral in HZV with skin involvement , will reduce post herpetic neuralgia



---

The cornea is protected against infection by:

- 1- Blinking
- 2- Washing of debris by the flow of tears
- 3- Entrapment of foreign particles by mucus
- 4- Anti-microbial properties of tear
- 5- The barrier function of the corneal epithelium

Loss of balance may predispose to infection



# Infective corneal lesions

## Bacterial Keratitis

---

- Over 90% of corneal infections are caused by bacteria
- Prime pathogens are staphylococcus and streptococcus
- Most bacteria are unable to penetrate the cornea if the epithelium is intact. (*Neisseria gonorrhoeae* is the only organism that can penetrate the intact epithelium)
  
- Predisposing factors:
  - a- Keratoconjunctivitis sicca (dry eye)
  - b- A breach in the corneal epithelium
  - c- Contact lens use
  - d- Prolonged use of topical steroids

---

## Symptoms:

- Pain
- Purulent discharge
- Ciliary injection
- Visual loss
- Hypopyon (white cells collected in the anterior chamber)



## Treatment:

- combined topical antibiotics to cover G+ve and G-ve
- Fluorquinolones can be used as monotherapy

# Infective corneal lesions

## Acanthameba Keratitis

Caused by a protozoa

- Common in contact lenses users



- Cause painful keratitis and ring shaped abscess

- Treated with chlorhexidene and propamidine for months and may end by corneal graft

# Infective corneal lesions

## Fungal Keratitis

- Common in warmer areas
- Should be considered in:
  - 1- Lack of response to antibacterial therapy
  - 2- Cases associated with prolong use of steroids
  - 3- History of trauma by plants → خصه زيتون  
دخل بالحيت
- Cause corneal opacity that appear fluffy.
- Treated with topical and systemic antifungals




# *Cornea*

---

## **Non- Infective corneal lesions**

# Corneal Dystrophies

- Rare inherited disorders, but relatively common in Jordan due to high consanguinity rates
- Affect corneal transparency
- Divided into:
  1. **Anterior Dystrophies** : present with recurrent corneal erosion
  2. **Stromal Dystrophies**: present with visual loss
  3. **Posterior Dystrophies** : gradual visual loss



# Disorders of Shape


## Keratoconus (KC)

القرنية  
المضروبة


Sporadic disorder associated with thinning of the centre of the cornea leads to ectasia and cone shaped cornea

Vision is affected without pain


\* the most important risk factor is eye rubbing ☆☆



Protrusion of the cornea cause myopia and astigmatism



# Sometimes mild or sublinical, Dx by corneal topography






# Modalities of treatment for KC

---


- Spectacles
- Contact lenses
- Corneal cross linking ●
- Intra stromal corneal rings ●
- Corneal graft ●

# Cross linking: stops progression of KC



increases bonds between collagen fibers. → stabilize it

# Rings reshape cornea and reduce astigmatism





# Corneal Grafting



---

- Donor corneal tissue can be grafted into a host cornea to restore corneal clarity or repair a perforation
- Avascular host cornea provides an immune-privileged site for grafting, with high success rate (80%)
- No need for HLA matching
- Extracted within 24 hours of death
- Topical steroids eye drops are used after operation to prevent graft rejection
- In Jordan, 50% of need is covered by local donation. JEB is based at JUH since 1979


**Cornea can be grafted within 2 weeks from extraction**

---





Non absorbable sutures for 1 year





# Corneal refractive surgery

---

- Principles
- Rate of success
- Contraindications


# Sclera ( white of the eye )

---

Composed of interwoven collagen fibrils of different widths

Variable in thickness, 1mm around optic nerve head and 0.3 mm posterior to muscle insertion

All six extraocular muscles are inserted on the sclera



# Diseases of the Sclera

---

**1- Episcleritis:** Inflammation of the superficial layer of the sclera

Causes mild discomfort with segmental redness of the eye

Usually self-limiting

If symptoms persist, topical anti-inflammatory treatment can be given

Rarely associated with systemic disease



## 2- Scleritis:

---

Usually associated with collagen vascular disease most commonly RA

Cause severe ocular pain with generalized redness of the eye

Characterized by swelling of the sclera






# Complications

---

- Scleromalacia (scleral thinning) sometimes with perforation
- Keratitis
- Uveitis
- Cataract formation
- Glaucoma



# Management

---

- Mild cases: topical steroids and oral NSAIDs
- Moderate and severe cases

Usually treated by high doses of systemic steroids or Cytotoxics

Medical condition that requires investigations for underlying cause



---

Thank you