


# *ENDOMETRIOSIS*

- The presence of a tissue similar to normal endometrium in structure and function outside the lining of the uterine cavity.
- Endom.interna → Adenomyosis
- Endom.externa → True endom.

# *ADENOMYOSIS*


## ■ Aetiology:

- Repeated Pregnancies.
- Vigorous Curettage.
- Hormonal Imbalance.




# *Adenomyosis---Pathology*

- **Symmetrical enlargement of uterus.**
- **Localized or diffuse.**
- **Histology:**
  - Glands +Stroma surrounded by muscle fibres.



# *Adenomyosis--Clinical features*

- End of reproductive life.
- Multiparous.
- Asymptomatic.
- Menorrhagia:
  - Enlarg. of uterus.
  - ↑ Blood supply.
  - Impaired contractility.
  - Ass.endom.hyperplas.
- Dysmenorrhea & Dyspareunia.



## *Adenomyosis-----cont.*

### ■ Myoma vs Adenomyosis


-Rarely enlarg.uterus >12-14wks.

-Regular enlarg.of the uterus.

■ Treatment  TAH


# **ENDOMETRIOSIS**

- **Implantation Theory(sampson)**
- **Coelomic Metaplasia.**
- **Lymphatic&Vascular Dissemin.**



# *Endometriosis--Predisp.factors*

- Age → 4th decade.
- Reprod.history → delay 1st pregn.
- High Social class.
- Genetic → 7% of 1st degree relat.  
1% of unrelated control
- Auto-immune.



# *Endometriosis---Increase*

- **Better ability to recognise the disease.**
- **The growing number of laparoscopic procedure.**
- **Emergence of predisposing factors.**
- **Patients and physicians----more aware of the disease.**



# *Endometriosis---Pathology*

## ■ Macroscopic:

-Small black dots(powder burn)




Large cystic masses(chocolate cysts)


- Others—black, dark brown, bluish puckered lesions, nodules.
- Atypical lesions:
  - Red implants( petechial, vesicular, polypoid, red flame like)
  - Serous or clear vesicles.
- White plaques and scarring.


# **Endometriosis--Pathology**


## ■ **Microscopic**

- Endometrial glands.
- Stroma.
- Evidence of bleeding.




















# *Endom. --- Clinical presentation*

- Asymptomatic → 25%
- Pain → -The commonest  
-Pelvic pain, Dysm, Dysp.
- Menorrhagia
- Infertility
- Acute abdomen
- Intermittent pyrexia



# *Endometriosis--presentation*

- Suggestive of endom:
  - Pelvic tenderness
  - Fixed retroverted uterus.
  - Tender uterosacral ligament.
  - Enlarged ovaries.
- Deeply infiltrating nodules---most reliably detected when clinical exam performed during menstruation.

# *Endometriosis--Diagnosis*

- Symptomatology.
- Defenitive Diagnosis:
  - Laparoscopy
  - Histology



# *Endometriosis--Diagnosis*

## ■ *Laparoscopy:*

- Gold standard investigation.
- Specific time in the menstrual cycle
  - Insufficient evidence.
- Classification systems----**subjective & correlate poorly with pain symptoms**

# *Endometriosis---Histology*

- Is it necessary----controversial.
- Positive histology-----confirm.
- Negative histology----doesn't exclude.
- Histological confirmation of at least one lesion is ideal.
- Endometriomas > 3 cm and deep infiltrating disease----Histology.

# CA 125

- **May be elevated.**
- **Compared with laparoscopy----has no value as a diagnostic tool.**



# *Endometriosis & Infertility*

- 15% of infertile women → Endom.
- 40-60% of endom. → Infertility
- Mechanisms:
  - Adhesions
  - Dyspareunia
  - ↑ prostaglandins
  - Tubal motility
  - Folliculogen.
  - C.L function
  - ↑ Macrophages
  - LUF
  - ↑ prolactine


# *Endometriosis--Treatment*

## ■ Depends on:

- Severity of symp. -Prev.Rx.
- Age -Fertility expectation.

## ■ Types:

- Expectant -Surgical -Medical




# *Endometriosis---Medical Rx*

- Endom.goes into remission during pregnancy → Pseudopregnancy
- Endom.invariably disappears after menopause → Pseudomenopause
- Androgen causes regression of endometriosis → Androgen

# *Endometriosis--Medical Rx*

- **Combined pills.**
- **Progestogen.**
- **Testosterone.**
- **Danazol.**
- **Gestrinone.**
- **GnRh agonists**
- **Aromatase inhibitors**



# *DANAZOL*


- Isoxazole derivative of 17-alpha-ethinyltestosterone.
- Action:
  - Bind to SHBG → ↑ Free testost.
  - ↓ Synthesis of SHBG by the liver
  - Prevent medcyclic surge of FSH,LH
  - Inhibits several enzym. processes involved in ovarian steroidogenesis
  - ↓ Estrogen & ↑ Androgen

## *Danazol---Side Effects*

- Weight gain.
- Fluid retention.
- ↓ Breast size.
- Growth of facial hair.
- Emotion.lability
- Fatigue.
- Oily skin
- Atrophic vagin.
- Muscle cramps.
- Irrever.deepen. of voice.
- ↑ Choles. ↓ HDL
- Insuline resist.


# **DANAZOL-----cont.**

- **Rx for 6-9 months.**
- **Dose 200mg twice daily.**
- **Contraindications:**
  - Pregnancy    -Breast feeding
  - Severe hepatic,cardiac,renal dis.
  - Thromboembolism    -Porphyria
  - Androgen dependent tumours



## *Medical Rx-----cont.*

- **Gesrinone:(Trienic-19-Norsteroid)**
  - Inhibits midcyclic surge of FSH,LH
  - Same side effects as danazol.
  - Long 1/2 life(2.5-5mg twice weekly)
- **GnRh agonists:**
  - Menopausal symptoms.
  - Breakthrough bleeding.
  - Loss of bone Ca.






# *Medical Treatment---cont*

- Aromatase Inhibitors:(anastrozole,letrozole)
  - Aromatase— enzyme that catalyzes the final and the key step of estrogen production.
  - Decrease both peripheral and local estradiol production.
  - May be better at suppressing local estrogen formation in endometriotic tissues than GnRH  
More effective.
  - Combined with ovarian suppression.

**FIGURE 2. Mechanism of local estrogen and prostaglandin biosynthesis in endometriosis**



AA—arachidonic acid  
PGH<sub>2</sub>—prostaglandin H<sub>2</sub>  
PGE<sub>2</sub>—prostaglandin E<sub>2</sub>  
cAMP—cyclic adenosine monophosphate

In endometriotic tissue, COX-2 regulates a key step in PGE<sub>2</sub> formation. It catalyzes the conversion of arachidonic acid (AA) to PGH<sub>2</sub>, which is then converted to PGE<sub>2</sub>. PGE<sub>2</sub> is the most potent known inducer of aromatase activity via a cAMP-mediated pathway. Aromatase catalyzes the conversion of androgens to estrogens, and estrogen, in turn, induces COX-2 production in uterine endothelial cells. Thus, a positive feedback cycle favors continuous production of PGE<sub>2</sub> and estrogens in endometriosis.

# *Endometriosis--Surgical Rx*

## ■ **Radical:**

TAH+Removal of as much endom.  
tissue as possible+Bilat.oophorect.

## ■ **Conservative:**

- Division of adhesions,Tuboplasty---
- Presacral neuroectomy
- Laser uterine nerve ablation

## *Medical treatment of endom associated pain.*

- **Empirical treatment without definitive diagnosis----Appropriate.**
  - Adequate analgesia.
  - Progestogens
  - Combined oral contraceptives.

## *Medical RX---cont*

- Effectiveness of NSAIDs----**inconclusive evidence.**
- Suppression of ovarian function for 6 months----**reduce pain.**
- Symptom recurrence is common following medical treatment.
- Aromatase inhibitor---**may be effective.**
- LNG-IUS-----**reduce pain**

# *Surgical Rx of Endom-associated pain*

- **Ideal practice –diagnose and remove surgically.**
- **Ablation----reduce pain.**
- **LUNA-----Doesn't reduce pain.**
- **Can be reduced by removing the entire lesions in severe and deeply infiltrating disease.**
- **Preop & postop hormonal rx----insufficient evidence of benefit.**

# *Treatment of Endom-associated Infertility*

- **Medical treatment:**
  - Minimal-mild disease----**Not effective and shouldn't be offered.**
  - More severe disease---**No evidence of effectiveness.**
- **Ablation & adhesiolysis----effective in minimal-mild disease.**
- **The role of surgery in improving pregnancy rate for moderate-severe disease is uncertain.**
- **Postop hormonal rx ---no beneficial effect.**

# *Assisted Reproduction in Endometriosis*

- IUI in minimal- mild---- Improves fertility.
- IVF is appropriate treatment:
  - Tubal function is compromised
  - Male factor
  - Other treatment have failed
- GnRH agonists for 3-6 months before IVF---  
increase rate of clinical pregnancy



Thank you